

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/99230/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

El-Hiti, Gamal A, Smith, Keith, Hegazy, Amany s. and Kariuki, Benson ORCID: <https://orcid.org/0000-0002-8658-3897> 2017. Crystal structure of 1,1-dimethyl-3-(4-methylphenyl)urea, C₁₀H₁₄N₂O. Zeitschrift für Kristallographie - New Crystal Structures 232 (2) , pp. 329-330. 10.1515/ncrs-2016-0290 file

Publishers page: <http://dx.doi.org/10.1515/ncrs-2016-0290>
<<http://dx.doi.org/10.1515/ncrs-2016-0290>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See


<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Open Access

Gamal A. El-Hiti*, Keith Smith, Amany S. Hegazy and Benson M. Kariuki

Crystal structure of 1,1-dimethyl-3-(4-methylphenyl)urea, $C_{10}H_{14}N_2O$



DOI 10.1515/ncrs-2016-0290

Received September 30, 2016; accepted January 13, 2017; available online February 21, 2017

Abstract

$C_{10}H_{14}N_2O$, monoclinic, *Cc* (no. 9), $a = 6.8354(9)$ Å, $b = 13.8221(17)$ Å, $c = 10.2905(12)$ Å, $\beta = 90.625(11)^\circ$, $V = 972.2(2)$ Å³, $Z = 4$, $R_{\text{gt}}(F) = 0.0526$, $wR_{\text{ref}}(F^2) = 0.1335$, $T = 296(2)$ K.

CCDC no.: 1527333

The asymmetric unit of the title crystal structure is shown in the figure. Tables 1 and 2 contain details on crystal structure and measurement conditions and a list of the atoms including atomic coordinates and displacement parameters.

Source of material

The title compound was synthesised by reaction of equimolar quantities of 4-methylaniline and dimethylcarbamoyl chloride in dichloromethane at 40 °C for 1 h in the presence of excess triethylamine (1.38 mole equivalents). The crude product was recrystallized from a 1:3 mixture (by volume) of ethyl acetate and diethyl ether to give 1,1-dimethyl-3-(4-methylphenyl)urea (92%) as colourless crystals, Mp.: 141–142 °C (lit. 137–138 °C [1]; 152–153 °C [2]).

*Corresponding author: Gamal A. El-Hiti, Cornea Research Chair, Department of Optometry, College of Applied Medical Sciences, King Saud University, P.O. Box 10219, Riyadh 11433, Saudi Arabia, e-mail: gelhiti@ksu.edu.sa

Keith Smith, Amany S. Hegazy and Benson M. Kariuki: School of Chemistry, Cardiff University, Main Building, Park Place, Cardiff CF10 3AT, UK

Table 1: Data collection and handling.

Crystal:	Colourless plate
Size:	$0.45 \times 0.35 \times 0.10$ mm
Wavelength:	Mo $K\alpha$ radiation (0.71073 Å)
μ :	0.8 cm^{-1}
Diffractometer, scan mode:	SuperNova, ω -scans
$2\theta_{\text{max}}$, completeness:	59.6° , >99%
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$, R_{int} :	3791, 3791, 0.059
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 2943
$N(\text{param})_{\text{refined}}$:	120
Programs:	CrysAlis ^{PRO} [17], SHELX [18], PLATON [19]

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.0013(4)	0.6008(2)	0.5771(3)	0.0391(7)
C2	−0.1240(5)	0.6047(3)	0.4684(4)	0.0517(10)
H2	−0.0968	0.5687	0.3945	0.062*
C3	−0.2885(5)	0.6629(3)	0.4726(4)	0.0584(11)
H3	−0.3714	0.6651	0.4004	0.070*
C4	−0.3341(4)	0.7180(3)	0.5803(4)	0.0489(9)
C5	−0.2091(4)	0.7138(3)	0.6860(4)	0.0489(8)
H5	−0.2366	0.7502	0.7596	0.059*
C6	−0.0424(4)	0.6559(3)	0.6848(4)	0.0452(8)
H6	0.0402	0.6543	0.7571	0.054*
C7	−0.5184(5)	0.7786(3)	0.5819(6)	0.0691(13)
H7A	−0.6305	0.7372	0.5721	0.104*
H7B	−0.5158	0.8242	0.5117	0.104*
H7C	−0.5259	0.8126	0.6630	0.104*
C8	0.2587(4)	0.4929(2)	0.4877(4)	0.0392(7)
C9	0.5152(6)	0.3823(3)	0.4257(5)	0.0587(10)
H9A	0.4308	0.3757	0.3511	0.088*
H9B	0.5441	0.3195	0.4608	0.088*
H9C	0.6347	0.4134	0.4007	0.088*
C10	0.5180(5)	0.4470(4)	0.6474(5)	0.0591(11)
H10A	0.6537	0.4311	0.6369	0.089*
H10B	0.4598	0.4025	0.7072	0.089*
H10C	0.5068	0.5116	0.6805	0.089*
N1	0.1702(4)	0.5411(2)	0.5869(3)	0.0422(6)
H1	0.2219	0.5347	0.6629	0.051*
N2	0.4186(4)	0.4404(2)	0.5234(4)	0.0490(7)
O1	0.1970(4)	0.4948(2)	0.3756(3)	0.0590(8)

Experimental details

The crystal was twinned and the structure was refined using option HKL 5 of the SHELX program [18]. All hydrogen atoms were placed in calculated positions and refined using a riding model. Methyl C-H bonds were fixed at 0.96 Å, with displacement parameters 1.5 times $U_{eq}(C)$, and were allowed to spin about the C—C bond. Aromatic C-H distances were set to 0.93 Å and N-H to 0.86 Å and their U_{iso} set to 1.2 times the U_{eq} for the atoms to which they are bonded.

Discussion

Urea derivatives are important intermediates in medicinal chemistry [3–5]. Many convenient processes have been reported for the production of substituted ureas [6–11]. Aromatic ureas can be substituted efficiently *via* lithiation chemistry [12–16].

The asymmetric unit consists of one molecule of C₁₀H₁₄N₂O. The molecule is not planar as indicated by torsion angles C2—C1—N1—C8 = 13.1(5)° and N1—C8—N2—C10 = –12.2(5)°. N—H···O hydrogen bonds form chains of molecules along [001].

Acknowledgements: The project was supported by King Saud University, Deanship of Scientific Research, Research Chair.

References

- Hutchby, M.: *Novel Synthetic Chemistry of Ureas and Amides*. Springer Thesis, Springer-Verlag, Berlin, Heidelberg, (2013).
- Franz, R. A.; Applegath, F.; Morriss, F. V.; Breed, L. W.: The preparation of unsymmetrical ureas from carbon monoxide, sulfur, and amines. *J. Org. Chem.* **27** (1962) 4341–4346.
- Kozikowski, A. P.; Zhang, J.; Nan, F.; Petukhov, P.; Grajkowska, A. E.; Wroblewski, J. T.; Yamamoto, T.; Bzdega, T.; Wroblewska, B.; Neale, J. H.: Synthesis of urea-based inhibitors as active site probes of glutamate carboxypeptidase II: efficacy as analgesic agents. *J. Med. Chem.* **47** (2004) 1729–1738.
- dos Santos, L.; Lima, L. A.; Cechinel-Filho, V.; Correa, R.; de Campos Buzzi, F.; Nunes, R. J.: Synthesis of new 1-phenyl-3-{4-[(2*E*)-3-phenylprop-2-enoyl]phenyl}-thiourea and urea derivatives with anti-nociceptive activity. *Bioorg. Med. Chem.* **16** (2008) 8526–8534.
- Khan, K. M.; Saeed, S.; Ali, M.; Gohar, M.; Zhaid, J.; Khan, A.; Perveen, S.; Choudhary, M. I.: Unsymmetrically disubstituted urea derivatives: a potent class of antiglycating agents. *Bioorg. Med. Chem.* **17** (2009) 2447–2451.
- Artuso, E.; Degani, I.; Fochi, R.; Magistris, C.: Preparation of mono-, di-, and trisubstituted ureas by carbonylation of aliphatic amines with *S,S*-dimethyl dithiocarbonate. *Synthesis* **39** (2007) 3497–3506.
- Mizuno, T.; Nakai, T.; Mihara, M.: Synthesis of unsymmetrical ureas by sulfur-assisted carbonylation with carbon monoxide and oxidation with molecular oxygen under mild conditions. *Synthesis* **41** (2009) 2492–2496.
- Wu, C.; Cheng, H.; Liu, R.; Wang, Q.; Hao, Y.; Yu, Y.; Zhao, F.: Synthesis of urea derivatives from amines and CO₂ in the absence of catalyst and. *Solvent. Green Chem.* **12** (2010) 1811–1816.
- Vinogradova, E. V.; Fors, B. P.; Buchwald, S. L.: Palladium-catalyzed cross-coupling of aryl chlorides and triflates with sodium cyanate: a practical synthesis of unsymmetrical ureas. *J. Am. Chem. Soc.* **134** (2012) 11132–11135.
- Carnaroglio, D.; Martina, K.; Palmisano, G.; Penoni, A.; Domini, C.; Cravotto, G.: One-pot sequential synthesis of isocyanates and urea derivatives via a microwave-assisted Staudinger–aza-Wittig reaction. *Beilstein J. Org. Chem.* **9** (2013) 2378–2386.
- Thalluri, K.; Manne, S. R.; Dev, D.; Mandal, B.: Ethyl 2-cyano-2-(4-nitrophenylsulfonyloxyimino)acetate-mediated Lossen rearrangement: single-pot racemization-free synthesis of hydroxamic acids and ureas from carboxylic acids. *J. Org. Chem.* **79** (2014) 3765–3775.
- Smith, K.; El-Hiti, G. A.; Alshammari, M. B.: Directed lithiation of *N'*-(2-(4-methoxyphenyl)ethyl)-*N,N*-dimethylurea and *tert*-butyl 2-(4-methoxyphenyl)ethyl)carbamate. *Synthesis* **46** (2014) 394–402.
- Smith, K.; El-Hiti, G. A.; Alshammari, M. B.: Control of site of lithiation of 3-(aminomethyl)pyridine derivatives. *Synthesis* **45** (2013) 3426–3434.
- Smith, K.; El-Hiti, G. A.; Alshammari, M. B.: Variation in the site of lithiation of 2-(2-methylphenyl)ethanamine derivatives. *J. Org. Chem.* **77** (2012) 11210–11215.
- Smith, K.; El-Hiti, G. A.; Alshammari, M. B.: Lithiation and substitution of *N'*-(ω-phenylalkyl)-*N,N*-dimethylureas. *Synthesis* **44** (2012) 20139–2022.
- Smith, K.; El-Hiti, G. A.; Hegazy, A. S.: Lateral lithiation of *N'*-(2-methylbenzyl)-*N,N*-dimethylurea and *N*-(2-methylbenzyl)pivalamide: Synthesis of tetrahydroisoquinolines. *Synthesis* **42** (2010) 1371–1380.
- Agilent Technologies: CrysAlis^{PRO} Software system, Agilent Technologies, Yarnton, England (2014).
- Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr. A* **64** (2008) 112–122.
- Farrugia, L. J.: WinGX and ORTEP for Windows: an update. *J. Appl. Crystallogr.* **45** (2012) 849–854.